

# Tick Bites



## Did You Know?

Lyme disease is carried by the deer tick or western black-legged tick. These ticks are harder to detect than dog ticks because they're much smaller (an adult tick is about the size of a sesame seed).

It's not uncommon to find a tick on a child. While most tick bites are harmless and don't require medical treatment, some ticks do carry harmful germs.

## What to Do:

1. If the tick is still attached to the child's skin, remove it:
  - Using fine-tipped tweezers, grasp the head of the tick close to the skin.
  - Firmly and steadily pull the tick straight out of the skin. **Do not** twist the tick, or rock it from side to side while removing it.
2. Put the tick in alcohol to kill it.
3. Wash your hands and the site of the bite with soap and water.
4. Swab the bite with alcohol.

## Call the child's doctor if:

- the tick may have been on the skin for more than 24 hours
- part of the tick remains in the skin after attempted removal
- the child develops a rash of any kind (especially a red-ringed bull's-eye rash)
- the area looks infected (increasing redness, warmth, swelling, pain, or oozing pus)
- the child develops symptoms like fever, headache, fatigue, chills, stiff neck or back, or muscle or joint aches

## Think Prevention!

When playing in wooded areas, children should wear long-sleeved shirts and pants. Spray insect repellent (containing no more than 10% to 30% DEET) on exposed skin and clothing. After kids play outside, check their skin – especially the scalp, behind the ears, the neck, under the arms, and the groin.

Note: All information is for educational purposes only. For specific medical advice, diagnoses, and treatment, consult your child's doctor. Review this with a doctor prior to use.